

Lepidii Semen



Figure 1 A photograph of Lepidii Semen

A. Lepidii Semen B. Magnified seed

Lepidii Semen**1. NAMES**

Official Name: *Lepidii Semen*

Chinese Name: 北葶藶子

Chinese Phonetic Name: Beitinglizi

2. SOURCE

Lepidii Semen is the dried ripe seed of *Lepidium apetalum* Willd. (Brassicaceae). The fruit is collected in summer when ripe, dried under the sun, then the seeds gathered and foreign matter removed to obtain *Lepidii Semen*.

3. DESCRIPTION

Ovoid and slightly flattened, 0.9-1.2 mm long, 0.6-0.9 mm wide. Externally brown to reddish-brown, slightly lustrous, marked with numerous petty granular protuberances and 2 longitudinal shallow furrows, one of which is more distinct than the other. One end obtuse, the other acuminate and slightly concave with indistinct hilum. Odour slight; taste slightly pungent and relatively viscous (Fig. 1).

4. IDENTIFICATION**4.1 Microscopic Identification** (*Appendix III*)**Transverse section**

Outer epidermis of testa consists of 1 layer of mucilage cells, with thickened inner-walls, extending outwards to cellulose columns. Cellulose columns oblique, lengthened, mucilage striations visible in periphery. Inner epidermis of testa consists of 1 layer of subsquare cells, with inner and lateral wall thickened and strongly lignified. Endosperm cells 1 layer, flat square, containing aleurone granules. Cotyledon composed of 2 slices, together with radicle occupying the major portion of the seed (Fig. 2).

Powder

Colour yellowish-brown. Outer epidermal cells of testa consist of 1 layer of mucilage cells, colourless or pale yellow, rectangular in lateral view, inner walls thickened, extending outwards to form cellulose column, cellulose column 15-46 μm long, apex oblique, lengthened or truncate, mucilage striations visible in periphery; rounded-polygonal in surface view, 25-56 μm in diameter, cellulose column visible at centre, mucilage striations visible in periphery. Inner epidermal cells of testa 1 layer in lateral view, yellow or reddish-brown, lateral walls unevenly thickened; polygonal, subsquare, sometimes long polygonal in surface view, 14-40 μm in diameter, walls 2-6 μm thick, lumens subrounded to polygonal. Endosperm cells colourless, flat square in section view, polygonal in surface view, walls slightly thickened, containing aleurone grains and oil droplets. Cotyledon cells irregular polygonal (Fig. 3).






Figure 2 Microscopic features of transverse section of *Lepidii Semen*.

A. Sketch B. Section illustration C. Testa magnified

- 1. Outer epidermis of testa 2. Cellulose column
- 3. Inner epidermis of testa 4. Endosperm 5. Cotyledons 6. Radicle




1-1



1-2




2-1



2-2



3



4

50 μm

Figure 3 Microscopic features of powder of Lepidii Semen (under the light microscope)

1. Outer epidermal cells of testa (1-1 in lateral view, 1-2 in surface view)
2. Inner epidermal cells of testa (2-1 in lateral view, 2-2 in surface view)
3. Endosperm cells 4. Cotyledon cells

4.2 Thin-Layer Chromatographic Identification [Appendix IV(A)]

Standard solution

Sinapic acid standard solution

Weigh 1.0 mg of sinapic acid CRS (Fig. 4) and dissolve in 10 mL of methanol.

Developing solvent system

Prepare a mixture of *n*-hexane, ethyl acetate and glacial acetic acid (10:6:1, v/v).

Test solution

Weigh 2.0 g of the powdered sample and place it in a 50-mL conical flask, then add 20 mL of methanol. Sonicate (220 W) the mixture for 30 min. Filter and transfer the filtrate to a 250-mL round-bottomed flask. Evaporate the solvent to dryness at reduced pressure in a rotary evaporator. Dissolve the residue in 1 mL of methanol.

Procedure

Carry out the method by using a HPTLC silica gel F₂₅₄ plate and a freshly prepared developing solvent system as described above. Apply separately sinapic acid standard solution (3 μL) and the test solution (4 μL) to the plate. Develop over a path of about 8 cm. After the development, remove the plate from the chamber, mark the solvent front and dry in air. Examine the plate under UV light (366 nm). Calculate the *R_f* value by using the equation as indicated in Appendix IV (A).




Figure 4 Chemical structures of (i) quercetin and (ii) sinapic acid




Figure 5 A reference HPTLC chromatogram of *Lepidii Semen* extract observed under UV light (366 nm)

1. Sinapic acid standard solution 2. Test solution

For positive identification, the sample must give spot or band with chromatographic characteristics, including the colour and the R_f value, corresponding to that of sinapic acid (Fig. 5).

4.3 High-Performance Liquid Chromatographic Fingerprinting (*Appendix XII*)

Standard solution

Sinapic acid standard solution for fingerprinting, Std-FP (50 mg/L)

Weigh 0.5 mg of sinapic acid CRS and dissolve in 10 mL of methanol (50%).

Test solution

Weigh 1.0 g of the powdered sample and place it in a 50-mL conical flask, then add 20 mL of methanol (50%). Sonicate (180 W) the mixture for 30 min. Filter and transfer the filtrate to a 25-mL volumetric flask. Make up to the mark with methanol (50%). Filter through a 0.45- μ m PTFE filter.

Chromatographic system

The liquid chromatograph is equipped with a DAD (324 nm) and a column (4.6 × 250 mm) packed with ODS bonded silica gel (5 μm particle size). The flow rate is about 1.0 mL/min. The mobile phase is a mixture of 0.1% acetic acid and acetonitrile (87:13, v/v). The elution time is about 60 min.

System suitability requirements

Perform at least five replicate injections, each using 10 μL of sinapic acid Std-FP. The requirements of the system suitability parameters are as follows: the RSD of the peak area of sinapic acid should not be more than 5.0%; the RSD of the retention time of sinapic acid peak should not be more than 2.0%; the column efficiency determined from sinapic acid peak should not be less than 10000 theoretical plates.

The *R* value between peak 2 and the closest peak in the chromatogram of the test solution should not be less than 1.0 (Fig. 6).

Procedure

Separately inject sinapic acid Std-FP and the test solution (10 μL each) into the HPLC system and record the chromatograms. Measure the retention time of sinapic acid peak in the chromatogram of sinapic acid Std-FP and the retention times of the three characteristic peaks (Fig. 6) in the chromatogram of the test solution. Identify sinapic acid peak in the chromatogram of the test solution by comparing its retention time with that in the chromatogram of sinapic acid Std-FP. The retention times of sinapic acid peaks from the two chromatograms should not differ by more than 2.0%. Calculate the RRTs of the characteristic peaks by using the equation as indicated in Appendix XII.

The RRTs and acceptable ranges of the three characteristic peaks of Lepidii Semen extract are listed in Table 1.

Table 1 The RRTs and acceptable ranges of the three characteristic peaks of Lepidii Semen extract

Peak No.	RRT	Acceptable Range
1	0.74	± 0.03
2 (marker, sinapic acid)	1.00	-
3	1.15	± 0.03




Figure 6 A reference fingerprint chromatogram of *Lepidii Semen* extract

For positive identification, the sample must give the above three characteristic peaks with RRTs falling within the acceptable range of the corresponding peaks in the reference fingerprint chromatogram (Fig. 6).

5. TESTS

5.1 Heavy Metals (*Appendix V*): meet the requirements.

5.2 Pesticide Residues (*Appendix VI*): meet the requirements.

5.3 Mycotoxins (*Appendix VII*): meet the requirements.

5.4 Sulphur Dioxide Residues (*Appendix XVII*): meet the requirements.

5.5 Foreign Matter (*Appendix VIII*): not more than 7.0%.

5.6 Ash (*Appendix IX*)

Total ash: not more than 7.5%.

Acid-insoluble ash: not more than 3.0%.

5.7 Water Content (*Appendix X*)

Oven dried method: not more than 9.0%.

6. EXTRACTIVES (*Appendix XI*)

Water-soluble extractives (cold extraction method): not less than 8.0%.

Ethanol-soluble extractives (cold extraction method): not less than 10.0%.

7. ASSAY

Carry out the method as directed in Appendix IV (B).

Standard solution

Quercetin standard stock solution, Std-Stock (100 mg/L)

Weigh accurately 0.5 mg of quercetin CRS (Fig. 4) and dissolve in 5 mL of methanol.

Quercetin standard solution for assay, Std-AS

Measure accurately the volume of the quercetin Std-Stock, dilute with methanol to produce a series of solutions of 1, 10, 20, 30, 40 mg/L for quercetin.

Test solution

Weigh accurately 1.0 g of the powdered sample and place it in a 100-mL round-bottomed flask, then add 30 mL of methanol and 0.5 mL of hydrochloric acid. Reflux the mixture for 2 h. Cool down to room temperature. Filter and transfer the filtrate to a 50-mL volumetric flask. Wash the residue for two times each with 10 mL of methanol. Filter and combine the filtrates. Make up to the mark with methanol. Filter through a 0.45- μ m PTFE filter.

Chromatographic system

The liquid chromatograph is equipped with a DAD (370 nm) and a column (4.6 \times 250 mm) packed with ODS bonded silica gel (5 μ m particle size). The flow rate is about 1.0 mL/min. The mobile phase is a mixture of 0.4% phosphoric acid and methanol (50:50, v/v). The elution time is about 25 min.

System suitability requirements

Perform at least five replicate injections, each using 10 μ L of quercetin Std-AS (20 mg/L). The requirements of the system suitability parameters are as follows: the RSD of the peak area of quercetin should not be more than 5.0%; the RSD of the retention time of quercetin peak should not be more than 2.0%; the column efficiency determined from quercetin peak should not be less than 5000 theoretical plates.

The *R* value between quercetin peak and the closest peak in the chromatogram of the test solution should not be less than 1.5.

Calibration curve

Inject a series of quercetin Std-AS (10 μ L each) into the HPLC system and record the chromatograms. Plot the peak areas of quercetin against the corresponding concentrations of quercetin Std-AS. Obtain the slope, y-intercept and the r^2 value from the 5-point calibration curve.

Procedure

Inject 10 µL of the test solution into the HPLC system and record the chromatogram. Identify quercetin peak in the chromatogram of the test solution by comparing its retention time with that in the chromatogram of quercetin Std-AS. The retention times of quercetin peaks from the two chromatograms should not differ by more than 5.0%. Measure the peak area and calculate the concentration (in milligram per litre) of quercetin in the test solution, and calculate the percentage content of quercetin in the sample by using the equations as indicated in Appendix IV (B).

Limits

The sample contains not less than 0.10% of quercetin ($C_{15}H_{10}O_7$), calculated with reference to the dried substance.